

Optical Monitoring of NLS1s at the Wise Observatory

Ohad Shemmer and Hagai Netzer

*School of Physics and Astronomy and the Wise Observatory, Tel-Aviv University,
Tel-Aviv 69978, Israel*

Abstract

We report the preliminary results of a spectrophotometric monitoring program of Narrow-Line Seyfert 1 galaxies conducted at the Wise Observatory in 1999. The data we collected will be used to evaluate the line-to-continuum time lag in an attempt to test the idea that the accretion rate in NLS1s is larger than in other Seyfert 1 galaxies.

Key words: galaxies: active; X-rays: galaxies

1 Observations and Preliminary Results

A spectrophotometric monitoring program of 5 Narrow-Line Seyfert 1 (NLS1) galaxies was conducted at the Wise Observatory (WO) during the summer and fall of 1999. The observations were performed with the WO spectrograph, and on separate occasions a CCD camera was used for broad-band photometry. Long-slit spectroscopy was carried out using a 10" wide slit for both the target NLS1s and nearby comparison stars, which were verified as non-variables by the photometric monitoring.

The prime target of observation was Akn 564, which was under a continuous monitoring campaign, coordinated by the AGN watch consortium, from both ground-based observatories and RXTE. In Figure 1 we present H α and optical continuum (observed 6950Å band) light curves for Akn 564, observed from WO. From the light curve it is evident that no significant variability was recorded for the object either on long or short time scales. In view of the extreme X-ray variability of NLS1s (preliminary results from the RXTE program show that during the first three months of the optical program, the object varied by over 100% in X-ray flux), this behavior is definitely surprising and is atypical of Seyfert 1 galaxies with broader emission lines.

Two more NLS1s from the sample did not show any significant variability: I Zw 1 and UCM 2257+2438, the latter (in sharp contrast with the former) showing very weak Fe II bands as opposed to other NLS1s (Figure 2). Mrk 335 and NGC 4051, which were also in our sample, varied on short as well as on long time scales, as already seen in previous monitoring campaigns.




Fig. 1. Akn 564. Light curves obtained from WO observations.




Fig. 2. The spectrum of UCM 2257+2438 obtained at the WO. Note the weak Fe II lines, unlike most NLS1s.